



# Cattle Producer's Handbook

Animal Health Section

667

## Bluetongue in Cattle

*Clell V. Bagley, D.V.M., Extension Veterinarian  
Utah State University*

Bluetongue (BT) is a viral disease that is spread mainly by one specific type of gnat. Other gnats and bloodsucking insects may occasionally transmit BT, but they are much less important in its transfer. Cattle are the main reservoir for overwintering the virus in temperate climates. The gnats become infected from cattle, and then spread the disease to other cattle and sheep as they take blood meals. Bluetongue is also spread through infected semen and may be spread by bloodsucking lice and a soft tick. It is possible that biting flies and contaminated needles may also spread it mechanically.

### Strains of Virus

There are 21 strains of BT virus that have been identified worldwide. Five are present in the United States: types 2, 10, 11, 13, and 17. The predominant strains that have been identified in the western U.S. are types 11, 13, and 17.

### Clinical Signs

Bluetongue is usually a much more serious disease in sheep than in cattle. A closely related virus causes epizootic hemorrhagic disease (EHD) in deer. The fatality rate is high for this virus in deer, but its effects on cattle and sheep are relatively mild.

There is still disagreement as to the extent of the clinical effects of BT in cattle. Some researchers feel it is a common cause of disease problems, and others say it is only rarely a problem. It is generally accepted that in the majority of infected adult cattle, the signs are inapparent.

There are considered to be three different clinical manifestations of BT in cattle:

1. Reproductive effects including abortion, infertility, mummification, and stillbirth.

2. Congenital defects with weak or dummy calves, deformed legs and feet, blindness, persistent covering of the gums over the front teeth, overshot lower jaw, and white-eye. The most susceptible period for the fetus to become infected is 60 to 140 days of pregnancy. Those affected later may still become latently (persistently) infected.
3. Persistently (latent) infected cattle with a hypersensitive reaction. These animals have apparently been previously infected with one strain of the virus but show no evidence until they are reinfected with it or another strain. The resulting disease condition may be moderately to extremely severe. Some of the signs identified include ulcers of the mouth, nose, esophagus, trachea, and rumen; lameness with an inflammation of the top of the hoof, laminitis, and hoof cracks; inflammation of the skin with loss of hair, sloughing of patches of skin, presence of an exudate and crusts on the skin surface, a burned muzzle, ulcers on the teats and udder; muscular stiffness; shallow rapid respiration with excessive salivation in stringy, long strands; and a swollen, protruding tongue. The temperature may be 106°F.

### Seasonality

Bluetongue is usually seen from midsummer until shortly after the first hard frost, when the gnats are inactivated. The congenital defects would not become evident until calving time. In areas with a mild winter climate, the gnats may be active year-round, and the disease could appear anytime.

### Diagnosis

Blood tests for antibodies are available for BT. A blood sample collected from a suspect affected calf be-